XN
A197
V.35 #2
V.38 #10

XC
0497
no.1-12

QK
0497
.11
1981

COMPOSITAE NEWSLETTER

LIBRARY
JUN 4 1981
NEW YORK
BOTANICAL GARDEN

Number Eleven February 1981

Charles Jeffrey, Editor, Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, U.K.

Financial support of the NEWSLETTER is generously provided by Otto Koeltz Antiquariat, P.O. Box 1360, 624 Koenigstein - Taunus, BRD, and by the Bentham-Moxon Trust, Royal Botanic Gardens, Kew.

NEWS FROM INDIVIDUALS

LANE, Meredith A., Assistant Professor, EPO Biology Dept., University of Colorado, 110 Ramaley, Campus Box B-334, Boulder , Colorado 80309, U.S.A., writes: My interests within the Compositae are primarily in the Astereae, as I have just completed my doctoral dissertation (at the University of Texas, under the direction of Prof. B. L. Turner) on the several genera Amphiachyris, Greenella, Gutierrezia, Gymnosperma, Thurovia and Xanthocephalum. At present, I am preparing the dissertation for publication in several parts, and am beginning an overview of the subtribe Solidagininae.

THANIKAIMONI, Dr. G., Institut Français, B.P. No. 33, Pondichéry - 605001, India, writes: I am getting several requests for the reprint of my paper "Principal works on pollen morphology of Compositae" published in "The Biology and Chemistry of the Compositae". Unfortunately, I have not received any reprint of this paper from the publishers. However, an updated checklist of 655 Asteraceae genera for which pollen morphological data is available is given in my "Fourth bibliographic index to the pollen morphology of Angiosperms" (Inst. fr. Pondichéry, trav.sec.sci.tech. Tome 17 : 1-338 : 1980), copies of which are available from the Director, French Institute, Pondicherry, India (Cost 125 FF + 10 FF "French Francs" including air mail charge).

EDITORIAL NOTE

I wish to point out to all recipients that the next Compositae Newsletter will be the last, unless an offer to edit, finance, produce and distribute the Newsletter is forthcoming within the next twelve months. I should be grateful to hear from anyone willing and able to undertake this service. Failing a previous solution, the International Botanical Congress in Sydney in August-September this year should provide an opportunity for interested synantherologists attending to discuss the future of the Newsletter. As I am not myself attending the Congress, I should be grateful to hear from a (preferably Australian) recipient willing to organise such a discussion.

C. Jeffrey

RECENT LITERATURE

These references have been extracted from the Current Awareness Lists of the Library, Royal Botanic Gardens, Kew. A comprehensive bibliography of Compositae, published annually, is contained in the Kew Record of Taxonomic Literature.

GENERAL

Barroso, G.M.
Levantamento das Compostas de Pernambuco.
In Sociedade Botânica do Brasil. XXXI Congresso Nacional de Botânica, 20-27 de janeiro de 1980. Itabuna, Ilhéus. 53-54 (1980). Abstract only.
Berhaut, J.
Flore illustrée du Sénégal. Dycotylédones: tome 2. Balanophoracées à Composées. Dakar, Gouvernement du Sénégal, Ministère du Développement Rural, Direction des Eaux et Forêts, 1974. 695 p. - Illus. Compositae p.418-663.
Cronquist, A.
Vascular flora of the southeastern United States: volume 1. Asteraceae. Chapel Hill, University of North Carolina Press, 1980. xv, 261p. - Chrom.nos., keys.
Espinar, L.A.
Notas sinonimicas sobre Compuestas Argentinas: 2.
Darwiniana, 22(4): 551-555 (1980). Eupatorium, Verbesina.
Grierson, A.T.C.
Compositae.
In Dassanayake, M.D., ed. A revised handbook to the flora of Ceylon: vol. 1. New Delhi, Amerind Publishing Co. Pvt. Ltd. 111-278 (1980) - En; chrom.nos., keys.
Jansen, R.K., Steussy, T.F.
Chromosome counts of Compositae from Latin America.
Amer. J. Bot., 67(4): 585-594 (1980).
Karim, F.M., Ali, H.H.
Pollen morphology in some species of Compositae.
Abu-Ghraib, Botany Directorate, 1979. [unpaged] - Illus.
Koster, J.T.
The Compositae of New Guinea: 7.
Blumea, 26(1): 233-243 (1980). - Illus., keys. Incl. Senecioneae.
Ling, Y. ed.
(Flora Reipublicae Popularis Sinicae: 75. Compositae (2): Inuleae-Helenieae).
Peking, Academia Sinica, 1979. 422p. - Ch; illus., keys.
Melhem, T.S.A., Silvestre, M.S.F., Makino, H.
Grãos de polen de plantas alerggénas: Compositae.
Hoehnea, 8:73-100 (1980) - (En); illus.

LACTUCEAE

Arevschatjan, I.G.
On Taraxacum stevenii DC. (De specie Taraxacum stevenii DC.)
Novit. Syst. Pl. Vasc., 8:282-283 (1978). Transl. from Novosti Sist. Vyssh. Rast., 8 (1971)
Bachman, K, Chambe s, K.L.
Genes regulat ng the appearance of two kinds of fruit in Microseris strain B87 (Asteraceae, Compositae).
Experientia 37: 29-31 (1981).
Bräutigam, S., Hilbig, W.
Ein neues Hieracium aus der Mongolischen Volksrepublik.
Ergebnisse der Mongolisch-Deutschen Biologischen Expeditionen seit 1962 Nr. 87.
Feddes Repert., 91(5-6): 273-280 - (En); illus., maps.
Chinnappa, C.C., Morton, J.K.
Unusual chromosome association in Krigia cespitosa (Compositae).
Caryologia, 32(4): 413-418 (1979 publ. 1980) - Illus., chrom. no.
Doll, R.
Neue Taraxaca aus Mecklenburg; 1.
Gleditschia, 5: 45-51 (1977) - Illus., map.
González, G.L.
Launaea fragilis (Asso) Pau, nombre correcto para L. resedifolia auct. plur. non (L.) Kuntze. (Launaea fragilis (Asso) Pau, the correct name for L. resedifolia auct. plur. non (L.) Kuntze.
An. Inst. Bot. A.J. Cavanilles, 36: 135-138 (1979) - (En).

LACTUCEAE (CONTINUED)

Hänsel, R., Kartarahardja, M., Huang, J.-T., Bohlmann, F.
Sesquiterenlacton-β-D-glucopyranoside sowie ein neues eudesmanolid aus Taraxacum officinale.
Phytochemistry, 19(5): 857-861 (1980). - (En).

Harriman, N.A.
Leontodon and Hypochaeris (Compositae) in Wisconsin.
Mich. Bot., 19(2): 93-95 (1980).

Jasiewicz, A.
Hieracium L., Jastrzebiec.
In Jasiewicz, A., ed. Flora polska: rośliny naczyniowe polski i ziem ościennych: tom. 14. Warszawa, Państwowe Wydawnicauto Naukowe, 199-335 (1980) - Pol; illus., chrom. no., key.

Jehlik, V.
Die Verhreitung von Lactuca tartarica in der Tschechoslowakei und Bemerkungen zu ihrem Vorkommen. (Rozsireni druhu Lactuca tartarica v Ceskoslovensku a poznámky k jeho výskytu.)
Preslia, 52(3): 209-216 (1980) - (En, Cz); map.

Lack, H.W., Ern, H., Straka, H.
Die Gattung Rothmaleria Font Quer (Asteraceae, Lactuceae).
Willdenowia, 10(1): 37-49 (1980) - (En); illus., map.

Lambert, D.S.
Taraxacum.
Irish Nat. J., 20(4): 169 (1980) - En. Records based on voucher specimens in the Ulster Museum.

Lehto, J.
"Extinct" wire - lettuce, Stephanomeria schotii (Compositae), rediscovered in Arizona after more than one hundred years.
Desert Pl., 1(1): 22 (1979).

Musina, L. S.
Klassifikatsiya vozmozhnye puti évolyutsii zhiznennykh form v podtribe Leontodontinae O. Hoffm. (Asteraceae). (Classification and possible paths of evolution of life forms of subtribe Leontodontinae O. Hoffm. (Asteraceae))
Byull. Mosk. Obshch. Ispyt. Prir., Biol., 85(2): 98-103 (1980) - (En).

Price, H.J., Bachmann, K., Chambers, K.L.
Intraspecific variation in genomic DNA content in Microseris.
Bot. Soc. Amer. Misc. Ser. Publ., 158: 91 (1980). Abstr. in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.

Rivas-Martinez, S., Saenz de Rivas, C.
Sobre Leontodon bourgaeanus Willk. (Asteraceae).
An. Inst. Bot. A.J. Cavanilles, 35: 155-157 (1978) - (En); map, keys.

Tacik, T.
Taraxacum Wiggers, Mniszek (Dmuchawiec).
In Jasiewicz, A., ed. Flora polska: rośliny naczyniowe polski i ziem ościennych: tom 14. Warszawa, Państwowe Wydawnictwo Naukowe. 7-199 (1980) - Pol; illus., chrom. nos., key.

Valentine, D.H.
Ecotypic and polymorphic variation in Centaurea scabiosa L.
Watsonia, 13(2): 103-109 (1980) - Illus.

Vasil'chenko, I.T.
What is Koelpinia latifolia C. Winkl.? (Quae est Koelpinia latifolia C. Winkl.?)
Novit. Syst. Pl. Vasc., 8: 277-281 (1978) - Illus., keys. Transl. from Novosti Sist. Vyssh. Rast., 8 (1971).

Watson, A.K.
The biology of Canadian weeds :43. Acroptilon (Centaurea) repens (L.) DC.
Canad. J. Pl. Sci. 60(3): 993-1004 (1980) - (Fr); illus., map, chrom. nos.

MUTISIEAE

Crisci, J.V.
Evolution in the subtribe Nassauviinae (Compositae, Mutisieae): a phylogenetic reconstruction.
Taxon, 29(2/3): 213-224 (1980) - Chrom. nos.

Golubkova, V.F.
New series in the genus Adenocaulon Hook. (Compositae). (Series novae generis Adenocaulon Hook. (Compositae))
Novit. Syst. Fl. Vasc., 9: 271 (1978). Transl. from Novosti Sist. Vyssh. Rast., 9 (1972).

Robinson, H.
Harthamnus, a new genus of Mutisieae from Bolivia (Asteraceae).
Phytologia, 45(6): 451-455 (1980) - Illus.

VERNONIEAE

Balbaa, S.I., Halim, A.F., Halaweish, F.T., Bohlmann, F.
New 5-methycoumarin derivatives from Ethulia conyzoides.
Phytochemistry, 19(7): 1519-1522 (1980).

Keeley, S.C., Jones, S.B., jr.
Distribution of pollen types in Vernonia (Vernonieae: Compositae).
Syst. Bot., 4(3): 195-202 (1979 publ. 1980) - Illus., map.

Leitão Filho, H. de, Semir, J.
Una nova combinação para o gênero Vernonia Schreb. (Compositae): Vernonia damazoi (Beauverd) Leitão Filho & Semir. (A new combination for the genus Vernonia Schreb. (Compositae).
Rev. Brasil. Bot., 2(2): 113-116 (1979) - (En); illus.

Robinson, H.
New species of Vernonieae (Asteraceae): 6. Lychnophora souzae.
Phytologia, 46(2): 104-106 (1980) - Illus.

Robinson, H.
New species of Vernonieae (Asteraceae): 7. Five new species of Vernonia from Brazil.
Phytologia, 46(2): 107-119 (1980) - Illus.

Robinson, H.
Re-establishment of the genus Critoniopsis (Vernonieae: Asteraceae)
Phytologia, 46(7): 437-442 (1980) - Illus.

Robinson, H.
New species of Vernonieae (Asteraceae): 7. Centratherum cardenasii from Bolivia.
Phytologia, 46(7): 443-445 (1980) - Illus.

Robinson, H., Bohlmann, F., King, R.M.
Chemosystematic notes on the Asteraceae: 3. Natural subdivisions of the Vernonieae.
Phytologia, 46(7): 421-436 (1980).

Rustaiyan, A., Nazarians, L., Bohlmann, F.
Two new 5-methycoumarins from Erlangea fusca.
Phytochemistry, 19(6): 1254-1255 (1980).

Stutts, J.G.
A new species of Vernonia (Compositae) from Ecuador.
Brittonia, 32(2): 152-153 (1980) - Illus.

LIABEAE

Robinson, H.
Flora of Ecuador no. 8. 190(2). Compositae - Liabeae.
Stockholm, Swedish Natural Science Research Council, 1978. 63p. - Illus., map, keys.

Robinson, H.
Studies in the Liabeae (Asteraceae): 14. New species of Liabum and Oligactis.
Phytologia, 46(2): 99-103 (1980) - Illus.

CARDUEAE

'Abd al-Hadi al Mukhtar, J.
Nabat al-Kassub. (The Kassub plant).
Bull. Dep. Bot. Min. Agric. Agrar. Ref. Rep. Iraq, no. 215. 18p. [n.d.] - Ara; illus.
Carthamnus.
Al-Sarraf, M.A.-A.A.
A taxonomic revision of the Iraqi species of Onopordum Linn. (Compositae), a thesis .... Master of Science in Botany. [Mosul], University of Mosul, 1979. ix, 93p. - (Pe); illus.
Bylova, A.M.
Bol'shoi zhiznennyi tsikl i vozrastnoi spektr tsenopopulyatsii vasil'ka sibirskogo. In Issled. sostava i izmenchiovosti flory i rastitel'nosti. M. 70-74 (1979). Abstr. in Ref. Zhurn., Biol., 7(2): 377 (1980).
Chen, Y.-L.
(Carlina L., a newly recorded genus of Compositae from China.)
Acta Phytotax. Sin., 18(2): 250, 255 (1980) - Ch; illus., key.
Couderc, M.
Quelques aspects du développement hétéroblastique dans le genre Crupina DC.
Bull. Soc. Bot. France, 126(3): 117-123 (1979) - (En); illus.
Dittrich, M., Petrak, F., Rechinger, K.H., Wagenitz, G.
Flora Iranica: no. 139b. Compositae: 3. Cynareae.
Graz, Akademische Druck, 1980. 287-468, 277-424p. - Illus., col. illus., keys.
Fernández Casas, J., Fernàndez Morales, M.J.
Centaurea lainzii, un triploide natural. (Centaurea lainzii, un triploide natural).
Mém. Soc. Bot. Genève, no. 1:115-122 (1979) - (Fr); illus., chrom. nos.
Georgiadis, T., Phitos, D.
Une Centaurea nouvelle du Mont Olympe (Grèce) de la section Phalolepis (Cass.) DC. Centaurea litochorea sp. nov.
Biol. Ecol. Medit., 5(1): 31-34 (1978) - (En); illus., chrom. nos.
Kamelin, R.V., Cherneva, O.V.
A new species of the genus Echinops L. from the section Chamaechinops Bunge. (Species nova generis Echinops L. e sectione Chamaechinops Bunge)
Novit. Syst. Pl. Vasc., 8: 262-265 (1978). Transl. from Novosti Sist. Vyssh. Rast., 8 (1971)
Lipshits, S. Yu
Understanding the genus Saussurea DC. :3. (Ad cognitionem generis Saussurea DC.:3)
Novit. Syst. Pl. Vasc., 8:266-271 (1978). Transl. from Novosti Sist. Vyssh. Rast., 8 (1971)
López, G.B.
Dos taxones nuevos del genero Centaurea L. (Two new taxa of the genus Centaurea L.)
An. Inst. Bot. A.J. Cavanilles, 36: 143-152 (1979) - (En); illus., map, chrom. nos.
López, G.B.
Notas cariosistematicas en el genero Centaurea L. Sect. Acrocentroides Willk. :1)
An. Inst. Bot. A.J. Cavanilles, 36: 349-369 (1979) - (En); illus., chrom. nos.
Ma, Y.-C.
(Revision of the genus Tugarinovia Iljin and its systematic position).
Acta Phytotax. Sin., 18(2): 217-219 (1980) - Ch (En); illus., key.
Mardaleĭshvili, T.K.
Ob odnom zabytom éndemichnom rastenii Kavkaza. (De una planta endemica florae Caucasi neglecta)
Zam. Sist. Geogr. Rast., 36: 58-63 (1980) - (Geo); illus. Psephellus buschiorum.
Moore, R.J.
The biology of Canadian weeds: 13. Cirsium arvense (L.) Scop.
In Mulligan, G.A., ed. The biology of Canadian weeds: contributions 1-32. Ottawa, Agriculture Canada, 146-161 (1979) - (Fr); illus., map, key. First publ. in Canad. J. Pl. Sci., 55: 1033-1048 (1975).
Podlech, D.
Jurinea merxmuelleri spec. nov. aus Afghanistan (Beiträge zur Flora von Afghanistan 12)
Mitt. Bot. Staatssamml. München, 16: 35-36 (1980).

CARDUEAE (CONTINUED)

Quinnell, R.J.
A taxometric survey of Arctium (Compositae) in Britain.
Unpubl., [London, St. Paul's School], n.d. 47, [16]p. - Illus., maps. Project for Biology A level.
Shtépa, I.S.
K voprosu defektnosti pŷl'tsŷv rode Cousinia Cass. (Ad cognitionem grana pollinis defectiva generis Cousinia Cass.)
Zam. Sist. Geogr. Rast., 36: 64-70 (1980) - (Geo).
Shin, C.
De genere Echinops L. familiae compositarum florae Sinicae.
Bull. Bot. Lab. North-east. Forest Inst., 5(5): 59-71 (1979). - Ch; illus.
Soskov, Yu D.
New nomenclature combination and series in the genus Rhaponticum Adans. (Combinatio nomenclaturica nova et series novae in genere Rhaponticum Adans.)
Novit. Syst. Pl. Vasc., 8: 275-276 (1978). Transl. from Novisti Sist. Vyssh. Rast.,8(1971)
Valdes-Bermejo, E., Rivas Goday, S.
Estudios en el genero Centaurea L. (Compositae). C. borjae sp. nov. (sect. Borjea sect. nov.)
An. Inst. Bt. A.J. Cavanilles, 35: 159-164 (1978) - (En); illus., chrom. nos., key.
Watson, A.K., Renney, A.J.
The biology of Canadian weeds: 6. Centaurea diffusa and C. maculosa.
In Mulligan, G.A., ed. The biology of Canadian weeds: contributions 1-32. Ottawa, Agriculture Canada. 62-76 (1979) - (Fr); illus., maps, chrom. no. First publ. in Canad. J. Pl. Sci.,54: 687-701 (1974).
Yano, K.
A new acetylenic alcohol from Cirsium japonicum.
Phytochemistry, 19(8): 1864-1866 (1980).

ARCTOTIDEAE

Friedrich, H.-C.
Arctotis merxmuelleri, eine neue Sippe aus dem kleinen Namaqualand.
Mitt. Bot. Staatssamml. München, 16:11-17 (1980) - Illus.

ANTHEMIDEAE

Barua, N.C., Sharma, R.P., Madhusudanan, K.P., Thyagarajan, G.,Herz, W.
Coumarins in Artemisia caruifolia.
Phytochemistry, 19(10) :2217-2218 (1980).
Biste, C.
Zytotaxonomische Untersuchungen des Formenkreises Achillea millefolium in der DDR.
Gleditschia, 6: 309 (1978) - Chrom. nos.
Borgen, L.
A new species of Argyranthemum (Compositae) from the Canary Island.
Norw. J. Bot., 27(3): 163-165 (1980) - Illus.
El-Emary, N.A., Bohlmann, F.
Zwei neue germacranolide aus Artemisia argentea.
Phytochemistry, 19(5): 845-847 (1980) - (En).
Elvira, R.
Plantas medicinales: .... manzanilla comun (Matricaria chamomilla L.)
Jardin Paisaje, 64(9): 21-23 (1980) - Illus.
Ilinskaya, A.F.
Porivnyal'no-anatomichne doslidzhennys epidermisu lŷstkiv deyakŷkh vŷdiv rodu Achillea L.(Comparative and anatomical studies of leaf epidermis in some Achillea L. species)
Ukr. Bot. Zhurn., 37(3): 25-29, 47 (1980) - (Rus, En); illus., chrom. nos.
Kelsey, R.G., Shafisadeh, F.
Glandular trichomes and sesquiterpene lactones of Artemisia nova.
Biochem. Syst. Ecol., 8(4): 371-377 (1980) - Illus.

ANTHEMIDEAE (CONTINUED)

Kovalevskaya, S.S.
New taxa of the genus Tanacetopsis Kovalevsk. (Taxa nova generis Tanacetopsis Kovalevsk.)
Novit. Syst. Pl. Vasc., 9: 272-274 (1978). Transl. from Novosti Sist. Vyssh. Rast., 9 (1972).

Kuvaev, V.B., Pakaln, D.A.
One variety of Taurian wormwood (Artemisia taurica Willd.) (Die varietate una Artemisiae tauricae Willd.)
Novit. Syst. Pl. Vasc., 8: 259-261 (1978) - Map. Transl. from Novosti Sist. Vyssh. Rast., 8 (1971).

Kuzmanov, B.A., Ninova, D.I., Kestenova, R. Kh.
Izmenchivost na Leucanthemum vulgare complex v Bulgariya : 2. Stravnitelno-morfologicho i anatomichno izledvane (1 chast). (Varibaility of Leucanthemum vulgare complex in Bulgaria: 2. Comparative morphological and anatomic investigation).
Fitologiya (Bulgaria), 15: 3-21 (1980) - (En); illus., key.

Lebedeva, T.S., Ochkivskaya, M.V.
Elektroforetychne vyvchennya bilkiv deyakykh bilokvitkovykh vydiv rodu Achillea L., shoho rostut' na Ukrayini. (Electrophoretic study of proteins in some albiflorous species from the Achillea L. genus growing in the Ukraine).
Ukr. Bot. Zhurn., 37 (5): 55-58 (1980).

Leonova, T.G.
O nekotorykh polynyakh (Artemisia L.) Mongol'skoi Narodnoi Respubliki. (De generis Artemisia L. speciebus nonnullis e Republica Populari Mogolica.)
Novosti Sist. Vyssh. Rast., 17: 233-239 (1980).

Leonova, T.G.
New species of wormwood (Artemisia L.) from the flora of Yukutian A.S.S.R. (Florae Jacutiae species novae e genere Artemisia L.)
Novit. Syst. Pl. Vasc., 8: 250-258 (1978) - Illus. Transl. from Novosti Sist. Vyssh. Rast., 8 (1971).

Ling, Y., Shih, C.
Taxa nova tribus Anthemidearum familiae Compositarum sinicarum.
Bull. Bot. Lab. North-East. Forest. Inst., no. 6: 1-16 (1980) - Ch. Brachanthemum, Dendranthema, Tripleurospermum, Pyrethrum, Cancrinia, Ajania.

Pireh, W., Tyul, R.J.
Cytogeography of Achillea millefolium in Oklahoma and adjacent states.
Rhodora, 82 (830): 361-367 (1980) - map, chrom. nos.

Redaelli, C., Formentini, L., Santaniello, E.
Apigenin 7-glucoside and its 2"- and 6"- acetates from ligulate flowers of Matricaria chamomilla.
Phytochemistry, 19(5): 985-986 (1980).

Toorn, J. van der
On the ecology of Cotula coronopifolia L. and Ranunculus sceleratus L. : 1. Geographic distribution, habitat, and field observations.
Acta Bot. Neerl., 29(5/6): 385-396 (1980) - Maps.

Tsvelev, N.N.
A new species of the genus Tanacetopsis (Tzvel.) Kovalevsk. from Central Asia. (Generis Tanacetopsis (Tzvel.) Kovalevsk. species nova ex Asia Media.)
Novit. Syst. Pl. Vasc., 8:248-249 (1978). Transl. from Novosti Sist. Vyssh. Rast., 8 (1971).

ASTEREAE

Anderson, L.C.
Occurrence and significance of glabrous achenes in Chrysothamnus (Asteraceae).
Bot. Soc. Amer. Misc. Ser. Publ., 158:7 (1980). Abstr. in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.

Anderson, L. C.
Morphology and biogeography of Chrysothamnus nauseosus ssp. texensis (Asteraceae): a new Guadalupe Mountains endemic.
Southwest. Nat., 25(2): 197-205 (1980) - Illus., chrom. no.

ASTEREAE (CONTINUED)

Beauchamp, R.M.
Baccharis vanessae, a new species from San Diego County, California.
Phytologia, 46(4): 216-222 (1980) - Illus., map.
Bohlmann, F., Robinson, H., King, R.M.
Chemotaxonomy of the genus Plagiocheilus.
Phytochemistry, 19(10): 2235-2236 (1980).
Brouillet, L.
Distribution of ploidy levels in Aster series Heterophylli: its significance.
Bot. Soc. Amer. Misc. Ser. Publ., 158: 17 (1980). Abstr. in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.
Clark, W.D., Brown, G.K., Mayes, R.A.
Pollen morphology of Haplopappus and related genera (Compositae - Astereae).
Amer. J. Bot., 67(9): 1391-1393 (1980) - Illus.
Clark, W.D., Urbatsch, L.E., Hartman, R.L., Mayes, R.A., Mabry, T.J.
Systematic implications of flavonoid patterns in Haplopappus segregates.
Biochem. Syst. Ecol., 8(3): 257-259 (1980) - Chrom. nos.
Espinar, L A.
Las especies centroargentinas de Hysterionica (Compositae).
Darwiniana, 22(4): 537-549 (1980) - (En); illus.
Given, D.R.
A taxonomic revision of Celmisia coriacea (Forst.f.) Hook.f. and its immediate allies (Astereae-Compositae).
New Zealand J. Bot., 18(1): 127-140 (1980) - Illus., maps, key.
Hill, L.M.
The genus Aster (Asteraceae) in Virginia.
Castanea, 45(2): 104-124 (1980) - Maps, key.
Jones, A.G.
A classification of the New World species of Aster (Asteraceae)
Brittonia, 32(2): 230-239 (1980) - Chrom. nos.
Jones, A.G.
Data on chromosome numbers in Aster (Astereae), with comments on the status and relationships of certain North American species.
Brittonia, 32(2): 240-261 (1980).
Lane, M.A.
New and re-instituted combinations in Gutierrezia (Compositae: Astereae).
Sida, 8(3): 313-314 (1980).
Melville, M., Morton, J.K.
A biosystematic study of the Solidago canadensis complex in Ontario.
Bot. Soc. Amer. Misc. Ser. Publ., 158: 74 (1980). Abstr. in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.
Schultz, L.M., Holmgren, A.H.
A new species of Townsendia (Asteraceae) from northern Arizona.
Brittonia, 32(2): 144-147 (1980) - Illus., key.
Semple, J.C.
A preliminary survey of flavonoids in tissue extracts of Aster sensu lato (Compositae)
Bot. Soc. Amer. Misc. Ser. Publ., 158: 101 (1980). Abstr. in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.
Semple, J.C., Brouillet, L.
A synopsis of North American Asters: the subgenera, sections and subsections of Aster and Lasallea.
Amer. J Bot., 67(7): 1010-1026 (1980) - Illus., chrom. nos., key.
Semple, J.C., Brouillet, L.
Chromosome numbers and satellite chromosome morphology in Aster and Lasallea.
Amer. J. Bot., 67(7): 1027-1039 (1980). - Illus.
Turner, B.L., Birdsong, A.
New combinations in the genus Aphanostephus (Astereae - Asteraceae)
Phytologia, 45(6): 501 (1980)

ASTEREAE (CONTINUED)

Urbatsch, L.E., Anderson, L.C.
Flavonoid survey of the genus Chrysothamnus (Asteraceae, Astereae).
Bot. Soc. Amer. Misc. Ser. Publ., 158:120 (1980). Abstr. in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.

CALENDULEAE

Praglowski, J., Grafström, E.
The pollen morphology of the tribe Calenduleae with reference to taxonomy.
Bot. Notiser, 133(2): 177-188 (1980) - Illus., key.

INULEAE

Arora, R.K., Maheshwari, M.L., Chandel, K.P.S., Gupta, R.
Mano (Inula racemosa):little known aromatic plant of Lahaul Valley, India.
Econ. Bot., 34(2): 175-180 (1980) - Illus.
Bohlmann, F., Abraham, W.-R., Scheldrick, W.S.
Weitere Diterpene mit Helifulvan-Gerüst und andere Inhaltsstoffe aus Helichrysum chionosphaerum.
Phytochemistry, 19(5): 869-871 (1980) - (En).
Bohlmann, F., Zdero, C., Abraham, W.R., Suwita, A., Grenz, M.
Neue Diterpene und neue Dihydrochalkon-Derivate sowie weitere Inhaltsstoffe aus Helichrysum - Arten.
Phytochemistry, 19(5): 873-879 (1980) - (En).
Bohlmann, F., Ziesche, J., King, R.M., Robinson, H.
Neue Eudesman-Derivate aus Pluchea suaveolens.
Phytochemistry, 19(5): 969-970 (1980).
Dakshini, K.M.M., Prithipalsingh
Numerical taxonomy of the genus Blumea in India.
Phytomorphology, 27(3): 247-260 (1977) - Illus.
Georgiadou, E., Lack, H.W., Merxmüller, H., Rechinger, K.H., Wagenitz, G.
Flora Iranica: no. 145. Compositae: 4. Inuleae.
Graz, Akademische Druck, 1980. 140, 128p. - Illus., keys.
Hilliard, O.M., Burtt, B.L.
Helichrysum fruticosum: its correct name and taxonomic status.
Notes Roy. Bot. Gard. Edinburgh, 38(1): 145-148 (1980).
Hilliard, O.M., Burtt, B.L.
(529) Proposal to conserve Petalacte D. Don (1826) against Billya Cassini (1825) (Compositae).
Taxon, 29(4): 507 (1980).
Jeffrey, C., Lack, H.W., Eldin, E.G., Wagenitz, G.
(539) Proposal to reject Pulicaria undulata (L.) C.A. Meyer under the terms of Article 69.
Taxon, 29(5/6): 694-695 (1980).
Nicoli, R.M., Buffard, Y.
Glanures corses: 10. Le genre Virginea et l'immortelle des frimas.
Bull. Soc. Linn. Provence, 31: 101-105 (1978) - (En).
Teresa, J.P., Barrero, A.F., San Feliciano, A., Medarde, M.
Eudesmane alcohols from Jasonia glutinosa.
Phytochemistry 19(10): 2155-2157 (1980).

HELIANTHEAE

Adsersen, H.
Revision of the Galápagos endemic genus Lecocarpus (Asteraceae).
Bot. Tiddsskr., 75(1): 63-76 (1980) - (Sp); illus., maps, chrom. nos., key.
Alonso, T.
El fichero de jardin: Tagetes patula.
Jard. Paisaje, 66(9): 43-44 (1980)- Col. illus.

HELIANTHEAE (CONTINUED)

Anaschenko, A.V.
Filogeneticheskie svyazi v rode Helianthus L. (Phylogenetic relations in the genus Helianthus L.)
Trudy Prikl. Bot.Genet. Selek., 64(2): 146-156 (1979) - (En).

Andrei, M., Moroşanu, E.
Die Entwicklung des männlichen Gametophyten von Helianthus annuus L. (Dezvoltarea gametofitului masculin la Helianthus annuus L.)
An. Univ. Bucuresti, Biol., 29: 3-8 (1980) - (Rum); illus.

Asai, Y.
(Additional notes on Verbesina alternifolia (L.) Britton in Japan.)
J. Jap. Bot., 55(7): 202-203 (1980) - Ja (En); illus.

Barsch-Gollnau, S., Ritterbusch, A., Mohr, H.
Photomorphogenesis and phyllotaxis during vegetative growth in Sinapis alba and Xanthium strumarium.
Pl., Cell, Environ., 3(5): 363-370 (1980) - Illus.

Bassett, I.J., Crompton, C.W.
The biology of Canadian weeds: 11. Ambrosia artemisiifolia L. and A. psilostachya DC.
In Mulligan, G.A., ed. The biology of Canadian weeds: contributions 1-32. Ottawa, Agriculture Canada. 120-133 (1979) - (Fr); illus., maps, chrom. nos. First publ. in Canad. J. Pl. Sci., 55:463-476 (1975)

Bate-Smith, E.C.
Astringent tannins of Cosmos bipinnatus.
Phytochemistry, 19(5): 982 (1980).

Becker, K.M.
A monograph of the genus Lasianthaea (Asteraceae)
Mem. New York Bot. Gard., 31(2): 1-64 (1979) - Illus., maps, chrom. nos., keys.

Best, K.F.
The biology of Canadian weeds: 10. Iva axillaris Pursh.
In Mulligan, G.A. ed. The biology of Canadian weeds: contributions 1-32. Ottawa, Agriculture Canada. 111-119 (1979) - (Fr); illus., map, chrom. nos., key. First publ. in Canad. J. Pl. Sci., 55: 293-301 (1975).

Bohlmann, F., Jakupovic, J., Dutta, L., Goodman, M.
Neue, abgewandelte Pseudoguajanolide aus Psilostrophe villosa.
Phytochemistry, 19(7): 1491-1494 (1980) - (En).

Bohlmann, F., Jakupovic, J., King, R.M., Robinson, H.
Neue Ent-Atisiren - und Ent - Kaurensäure - Derivate aus Helianthus - Arten.
Phytochemistry, 19(5): 863-868 (1980) - (En).

Bohlmann, F., Jakupovic, J., King, R.M., Robinson, H.
Chromones and flavans from Marshallia obovata.
Phytochemistry, 19(8): 1815-1820 (1980)

Bohlmann, F., Jakupovic, J., Robinson, H., King, R.M.
Neue Diterpene aus Schkuhria - Arten.
Phytochemistry, 19(5): 881-884 (1980) - (En).

Bohlmann, F., Knoll, K.-H., Robinson, H., King, R.M.
Neue Eudesmanolide aus Steiractinia mollis.
Phytochemistry, 19(5): 971-972 (1980).

Bohlmann, F., Zdero, C., Cuatrecas, J., King, R.M., Robinson, H.
Neue Sesquiterpene und Norditerpene aus Vertretern der Gattung Libanothamnus.
Phytochemistry, 19(6): 1145-1148 (1980) - (En).

Bohlmann, F., Zdero, C., King, R.M., Robinson, H.
New sesquiterpene lactones and other constituents from Fitchia speciosa.
Phytochemistry, 19(6): 1141-1143 (1980)

Bohlmann, F., Ziesche, J., King, R.M., Robinson, H.
Neue melampolide aus Smallanthus fruticosus.
Phytochemistry, 19(5):973-974 (1980).

Bohlmann, F., Ziesche, J., Robinson, H., King, R.M.
Neue amide aus Spilanthes alba.
Phytochemistry, 19(7): 1535-1537 (1980).

HELIANTHEAE (CONTINUED)

Cabrera, A.L.
Sobre la presencia del genero Arnica (Compositae) en la Argentina.
Darwiniana, 22 (4): 587-589 (1980).
Carr, G.D.
Two new species in the Hawaiian endemic genus Dubautia (Compositae)
Pac. Sci., 33(3): 233-237 (1979) - Illus., chrom. nos.
Clark, C., Thompson, W.C., Kyhos, D.W.
Comparative morphology of the leaf trichomes of Encelia (Compositae: Heliantheae)
Bot. Soc. Amer. Misc. Ser. Publ., 158:23 (1980). Abstract in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.
Crawford, D.J., Smith, E.B.
Flavonoid chemistry of Coreopsis grandiflora (Compositae)
Brittonia, 32(2): 154-159 (1980) - Chrom. nos.
Cuatrecasas, J.
Miscellaneous notes on neotropical flora: 12.
Phytologia, 47(1):1-13 (1980) - Chrom. no. Libanothamnus, Espeletia.
Domingues, X.A., Franco, R., Bohlmann, F., Bohlmann, R.
A pseudoguaianolide from Hymenoxys linearifolia.
Phytochemistry 19(10): 2204-2205 (1980)
Dunlop, C.R.
Dichromochlamys, a new genus in Asteraceae (Astereae).
J. Adelaide Bot. Gard., 2(3):235-239 (1980) - Illus., map.
Dunlop, C.R.
A revision of Ixiochlamys (Asteraceae: Astereae).
J. Adelaide Bot. Gard., 2(3): 241-252 (1980) - Illus., maps, key.
Funk, V.A., Raven, P.H.
Polyploidy in Montanoa Cerv. (Compositae, Heliantheae).
Taxon, 29(4): 417-419 (1980) - Illus., chrom. nos.
Gamisans, J.
A propos de Bellium nivale Req. (Notice on Bellium nivale Req.)
Candollea, 35(1): 201-209 (1980) - (En); chrom. nos.
Herz, W., Govindan, S.V.
Eucannabinolide and other constituents of Schkuhria virgata.
Phytochemistry, 19(6): 1234-1237 (1980)
Inouye, D.W., Inouye, R.S.
The amino acids of extrafloral nectar from Helianthella quinquenervis (Asteraceae).
Amer. J. Bot., 67(9): 1394-1396 (1980).
Kumamota, J., Scora, R.W.
Phytochemical investigations in Parthenium.
Bot. Soc. Amer. Misc. Ser. Publ., 158:61 (1980). Abstr. in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.
Lack, H.W.
The genus Enydra (Asteraceae, Heliantheae) in West Tropical Africa.
Willdenowia, 10(1): 3-12 (1980) - Illus., map, key.
Mears, J.A.
Flavonoid diversity and geographic endemism in Parthenium.
Biochem. Syst. Ecol., 8(4): 361-370 (1980).
Mooring, J.S.
A cytogeographic study of Chaenactis douglasii (Compositae, Helenieae).
Amer. J. Bot., 67(9): 1304-1319 (1980) - Map, chrom. nos.
Morley, B.
The nomenclature of and a key to some cultivated species of Montanoa Cervantes (Compositae).
J. Adelaide Bot. Gard., 2(2): 151-161 (1980) - Illus., key.

HELIANTHEAE (CONTINUED)

Neto, H.M., Guimarães, D.J., Marquete, O., Magalhaẽs, H.G., Oliveira, B.A.
Estudo geral de Stevia rebaudiana (Bertoni) Bertoni 1905:4. Diagnose e tipus. In Sociedade Botânica do Brasil, XXXI Congresso Nacional de Botąnica, 20 a 27 de janeiro de 1980. Itabuna, Ilhéus. 68 (1980). Abstract only.

Olsen, J., Mabry, T.J.
Systematic implications of flavonols in Zaluzania. Biochem. Syst. Ecol., 8(3): 261-262 (1980) - Chrom. nos.

Ortega, A., Lara, R., Martinez, R., Diaz, E.
Sphaerocephalin, a germacranolide isolated from Viguiera sphaerocephala. Phytochemistry, 19(7): 1545-1546 (1980).

Oswald, F.W.
A new form of Helianthus from south-west Florida. Phytologia, 44(6): 19-20 (1979).

Parker, K.F.
New combinations in Tetraneuris Greene (Heliantheae, Asteraceae). Phytologia, 45(6): 467 (1980)

Patil, T.M.
Phospholipid composition in Parthenium hysterophorus Linn. Geobios (Jodhpur), 7(4): 172-173 (1980).

Pieman, A.K., Towers, G.H.N., Rao, P.V.S.
Coronopilin - another major sesquiterpene lactone in Parthenium hysterophorus. Phytochemistry, 19(10): 2206-2207 (1980)

Roberts, M.L.
The flavonoids of Megalodonta beckii and isolation of 2'-3-Dihydroxy-4-methoxy-4'-glucosyl chalcone. Biochem. Syst. Ecol., 8(2): 115-118 (1980)

Robinson, H.
Studies in the Heliantheae (Asteraceae): 23. New Andean species Verbesina and Viguiera. Phytologia, 45(6): 456-459 (1980).

Robinson, H.
Studies in the Heliantheae (Asteraceae): 24. A new subtribe, Lycapsinae. Phytologia, 46(2): 120 (1980).

Robinson, H.
Studies in the Heliantheae (Asteraceae): 25. A new species of Unxia. Phytologia, 46(5): 289-292 (1980) - Illus.

Quijano, L., Olivier, E.J., Fischer, N.H.
Tetraludins D to N, eleven new melampolides from Tetragonotheca ludoviciana. Phytochemistry, 19(7): 1485-1489 (1980).

Seaman, F.C., Fischer, N.H.
New germacrolide- and heliangolide-type sesquiterpene lactones from Melampodium linearilobum. Phytochemistry, 19 (5): 849-856 (1980)

Seaman, F.C., Fischer, N.H., Steussy, T.F.
Systematic implications of sesquiterpene lactones in the subtribe Melampodiinae. Biochem. Syst. Ecol., 8(3): 263-271 (1980)

Schilling, E.E.
Flavonoid chemistry and the systematics of Helianthus (Compositae). Bot. Soc. Amer. Misc. Ser. Publ., 158:100 (1980). Abstr. in Botany 80: abstracts of papers to be presented at the University of British Columbia, Vancouver, 12-16 July 1980.

Schmitz, R., Frahm, A.W., Kating, H.
Ein neues Sesquiterpenketon aus Arnica - Arten. Phytochemistry, 19(7): 1477-1480 (1980) - (En).

Sørensen, P.D.
New taxa in the genus Dahlia (Asteraceae, Heliantheae - Coreopsidinae). Rhodora, 82(830): 353-360 (1980). - Illus., map, chrom. nos.

3 5185 00288 3831

HELIANTHEAE (CONTINUED)

Straley, G.B.
Arnica in the wild and in cultivation in British Columbia.
Davidsonia, II(2): 25-33 (1980) - Illus., key.
Turner, B.L., Davies, F.
Steussya (Asteraceae: Heliantheae), a new genus from south-central Mexico.
Brittonia, 32(2): 209-212 (1980) - Illus.
Turner, B.L., Dawson, D.
Taxonomy of Tetragonotheca (Asteraceae - Heliantheae).
Sida, 8(3): 296-303 (1980) - Illus., maps, chrom. nos., key.
Ulubelen, A., Kerr, K.M., Mabry, T.J.
New 6- hydroxyflavonoids and their methyl ethers and glycosides from Neurolaena oaxacana.
Phytochemistry, 19(8): 1761-1766 (1980).
Vivar, A.R. de, Bratoeff, E., Ontiveros, E., Lankin, D.C., Bhacca, N.S.
Viguilenin, a germacranolide from Viguiera linearis.
Phytochemistry, 19(8): 1795-1797 (1980).
Wolf, S.J.
Cytogeographical studies in the genus Arnica (Compositae: Senecioneae): 1.
Amer. J. Bot., 67(3): 300-308 (1980) - Chrom. nos.
Wussow, J.R., Urbatsch, L.E.
A systematic study of the genus Tetrachyron (Asteraceae: Heliantheae)
Syst. Bot., 4(4): 297-318 (1979) - Illus., map, key.

EUPATORIEAE

Bohlmann, F., Rosenberg, E., King, R.M., Robinson, H.
Neue Labdan-Derivate aus Aristeguietis buddleaefolia.
Phytochemistry, 19(5): 977-979 (1980) - (En).
Bohlmann, F., Suwita, A., King, R.M., Robinson, H.
Neue guajanolide aus Eupatorium rotundifolium.
Phytochemistry, 19(6): 1233-1234 (1980)
Bohlmann, F., Zdero, C., King, R.M., Robinson, H.
New heliangolides from Conocliniopsis prasiifolia.
Phytochemistry, 19(7): 1547-1549 (1980)
Bohlmann, F., Zdero, C., Robinson, H., King, R.M.
Guaianolides from Agrianthus pungens.
Phytochemistry, 19(8): 1873-1874 (1980).
King, R.M., Robinson, H.
Studies in the Eupatorieae (Asteraceae): 178. Additions to Eitenia and Lomatozoma.
Phytologia, 44(7): 455-462 (1979) - Illus., key.
King, R.M., Robinson, H.
Studies in the Eupatorieae (Asteraceae): 184. A new species of Arrojadocharis.
Phytologia, 44(7): 463-465 (1979) - Illus.
King, R.M., Robinson, H.
Studies in the Eupatorieae (Asteraceae): 185. Additions to the genus Lasiolaena.
Phytologia, 44(7): 466-472 (1979) - Illus., key.
King, R.M., Robinson, H.
Studies in the Eupatorieae (Asteraceae): 190. A new genus Santosia.
Phytologia, 45(6): 463-466 (1980).
King, R.M., Robinson, H.
Studies in the Eupatorieae (Asteraceae): 191. Various new species from Brasil.
Phytologia, 46(5): 295-307 (1980) - Illus. Morithamnus, Stevia, Lasiolaena, Koanophyllon, Disynaphia, Dasycondylus, Campuloclinium.
King, R.M., Robinson, H.
Studies in the Eupatorieae (Asteraceae): 192. Validation of subtribes.
Phytologia, 46(7): 446-450 (1980) - Chrom nos.
King, R.M. , Robinson, H., Barroso, G.M.
Studies in the Eupatorieae (Asteraceae): 177. A new genus, Morithamnus.
Phytologia, 44(7): 451-454 (1979) - Illus.

EUPATORIEAE (CONTINUED)

Ulubelen, A., Timmermann, B.N., Mabry, T.J.
Flavonoids from Brickellia chlorolepis and B. dentata.
Phytochemistry, 19(5): 905-908 (1980).

SENECIONEAE

Bohlmann, F., Dhar, A.K., Ahmed, M.
Thymol derivative from Doronicum hungaricum.
Phytochemistry, 19(8): 1850-1851 (1980).
Bohlmann, F., Ziesche, J.
New sesquiterpenes from Senecio species.
Phytochemistry, 19(8): 1851-1853 (1980)
Bohlmann, F., Zdero, C.
New furanoeremophilane derivatives from Ligularia thyrsoidea.
Phytochemistry, 19(7):1550-1551 (1980)
Cabrera, A.L., Zardini, E.M.
Sinopsis preliminar de las especies argentinas del género Senecio (Compositae)
Darwiniana, 22(4):427-492 (1980) - Key.
Davies, F.G.
The genus Gynura (Compositae) in India, Sri Lanka and the Seychelles.
Kew Bull., 35(2): 363-367 (1980) - Keys.
Lawrence, M.E.
Senecio L. (Asteraceae) in Australia: chromosome numbers and the occurrence of polyploidy.
Austral. J. Bot., 28(2): 151-165 (1980) - Illus.
Nordenstam, B.
A new species of Senecio from Namibia.
Mitt. Bot. Staatssamml. München, 16:29-33 (1980) - Illus.
Robinson, H.
Studies in the Senecioneae (Asteraceae): 10. A new species of Senecio from Brasil.
Phytologia, 46(5):293-294 (1980) - Illus.
Röder, E., Wiedenfeld, H., Frisse, M.
Pyrrolizidinalkeloide aus Senecio doronicum.
Phytochemistry, 19(6): 1275-1277 (1980).
Rundel, J.A., Rundel, P.W.
Succulent arborescent Senecios from Central Mexico.
Cact. Succ. J. (U.S.A.), 52(5):233-235 (1980) - Illus., col. illus., map.
Schmitt, J.
Pollinator foraging behavior and gene dispersal in Senecio (Compositae).
Evolution, 34(5): 934-943 (1980)
St. John, H.
The native species of Senecio (Compositae) in Hawaii. Hawaiian plant studies 53.
Pac. Sci., 33(4):329-332 (1979 publ. 1980) - Illus.
Weir, J., Ingram, R.
Ray morphology and cytological investigations of Senecio cambrensis Rosser.
New Phytologist, 86(2):237-241 (1980) - Chrom. nos.

Zdero, C., Bohlmann, F., Robinson, H., King, R.M.
Neue Furanoeremophilane aus Gynoxys dielsiana.
Phytochemistry, 19(5): 975-976 (1980).

N.B. Details of papers giving new floristic records for Britain, Canada (Québec), Germany (BRD), Iberian Peninsula, India (E), Patagonia, Spain, Sweden and U.S.S.R. (including Krŷm, Colchis, Middle Asia and Primorye), not cited above, can be obtained from the Editor.

Erratum: the two entries under Dunlop, C.R., and that under Gamisans, J., listed in Heliantheae, should be transferred to Astereae - Ed.

XN
A1958
V. 1 #3
V. 5 #1